

DISCUSSION

In microscopic polyangiitis, necrotising glomerulonephritis is the most common abnormality and pulmonary capillaritis often occurs. Perinuclear ANCA are present in more than 60% of patients, and antimyeloperoxidase antibody is closely allied.¹ Gastrointestinal involvement is usually mild and there is no report of extended colonic ulcerations and haemorrhage.² In addition, colonic biopsy disclosed crypt abscess, which can be found in ulcerative colitis, infectious colitis, radiation induced colitis, and graft versus host disease.³⁻⁵ However, such a finding has not been reported previously in patients with microscopic polyangiitis. Ulcerative colitis was diagnosed initially according to the colonoscopic and pathological findings. Pulmonary vasculitis is very rare in patients with inflammatory bowel diseases.⁶ Renal disease with necrotising glomerulonephritis is not commonly seen in patients with ulcerative colitis.⁷ In addition, antimyeloperoxidase antibody is usually not detected in such patients.⁸ The clinical course of this patient who had lung and renal manifestations did not favour a diagnosis of ulcerative colitis. The final diagnosis was microscopic polyangiitis with gastrointestinal involvement.

Extended colonic ulcerations with haemorrhage have not been reported previously in patients with microscopic polyangiitis. Vasculitis should be considered for differential diagnoses of colonic ulcerations, especially when the presentation is atypical, to avoid delay of prompt treatment.

Authors' affiliations

C-N Tsai, M-F Liu, C-R Wang, Section of Rheumatology, Department of Internal Medicine, College of Medicine, National Cheng Kung University, Tainan, Taiwan

C-M Chang, Section of Geriatrics, Department of Internal Medicine, College of Medicine, National Cheng Kung University, Tainan, Taiwan
 C-H Chuang, Section of Gastroenterology, Department of Internal Medicine, College of Medicine, National Cheng Kung University, Tainan, Taiwan
 Y-T Jin, Department of Pathology, College of Medicine, National Cheng-Kung University, Tainan, Taiwan

Correspondence to: Dr C-R Wang, Section of Rheumatology and Section of Allergy and Immunology, Department of Internal Medicine, College of Medicine, National Cheng Kung University, No 1, University Road, Tainan, Taiwan; wangcr@mail.ncku.edu.tw

Accepted 10 March 2004

REFERENCES

- Jennette JC, Falk RJ. Small-vessel vasculitis. *N Engl J Med* 1997;**337**:1512-23.
- Guillevin L, Durand-Gasselino B, Cevallos R, Gayraud M, Lhote F, Collard P, et al. Microscopic polyangiitis: clinical and laboratory findings in 85 patients. *Arthritis Rheum* 1999;**42**:421-30.
- Boyd JF. Pathology of the alimentary tract in Salmonella typhimurium food poisoning. *Gut* 1985;**26**:935-44.
- Northway MG, Scobey MW, Geisinger KR. Radiation proctitis in the rat. Sequential changes and effects of anti-inflammatory agents. *Cancer* 1988;**62**:1962-9.
- Epstein RJ, McDonald GB, Sale GE, Shulman HM, Thomas ED. The diagnostic accuracy of the rectal biopsy in acute graft-versus-host disease: a prospective study of thirteen patients. *Gastroenterology* 1980;**78**:764-71.
- Storch I, Sachar D, Katz S. Pulmonary manifestations of inflammatory bowel disease. *Inflamm Bowel Dis* 2003;**9**:104-15.
- Wilcox GM, Aretz HT, Roy MA, Roche JK. Glomerulonephritis associated with inflammatory bowel disease. *Gastroenterology* 1990;**98**:786-91.
- Abad E, Tural C, Mirapeix E, Cuxart A. Relationship between ANCA and clinical activity in inflammatory bowel disease: variation in prevalence of ANCA and evidence of heterogeneity. *J Autoimmun* 1997;**10**:175-80.

Severe digital ischaemia treated with phosphodiesterase inhibitors

C R Kumana, G T Y Cheung, C S Lau

Ann Rheum Dis 2004;**63**:1522-1524. doi: 10.1136/ard.2003.015677

Pulmonary hypertension is associated with autoimmune diseases, giving rise to digital ischaemia, including Raynaud's phenomenon (RP). As sildenafil, a phosphodiesterase-5 (PDE-5) inhibitor, relieves the pulmonary hypertension,¹ we suggested that it might also benefit digital ischaemia and RP symptoms. This report describes the impact of oral sildenafil (Viagra, donated by Pfizer HK) on three women with autoimmune disease and progressively severe digital ischaemia, despite treatment with a variety of drugs, including intravenous iloprost (fig 1). All patients gave written informed consent to try this off-label treatment.

CASE REPORTS

A 42 year old woman with dermatomyositis "sine myositis" and thyrotoxicosis (case 1) taking prednisolone, hydroxychloroquine, and carbimazole, experienced regression of troublesome RP and ischaemic digital ulceration after receiving diltiazem SR 200 mg/day and thrice monthly iloprost infusions. Ten months later, antituberculosis chemotherapy was started and the steroid dosage reduced, after she developed a febrile illness with cough, pleuropericardial effusions, and sputum culture positive for tuberculosis (TB). Digital gangrene supervened in both hands and both feet despite daily intravenous iloprost infusions for 2 weeks.

A day after starting sildenafil 50 mg three times a day her digital circulation and pain improved markedly; facial flushing was the only adverse reaction. In view of the slightly deteriorating digital ischaemia and experimental evidence,^{2,3} we inferred that rifampicin treatment was inducing the metabolism of sildenafil. Its dosage was therefore doubled. Symptomatic improvement continued and ischaemic tissues became demarcated. However, 26 days after starting sildenafil she succumbed to uncontrolled pulmonary TB.

A 28 year old woman with scleroderma/lupus (case 2) who had been receiving diltiazem, domperidone, omeprazole, and penicillamine for about 4 years presented with progressive breathlessness. Computed tomography of the thorax suggested fibrosing alveolitis; prednisolone and azathioprine were started. Thereafter, she incurred occasional chest infections, other complications, and features of autoimmune diseases. She then developed persistent high grade fever, chills, rigors, night sweats, and increasingly severe digital arthralgia, vasculitis, and ischaemia. A polymerase chain reaction (PCR) of blood disclosed disseminated TB. Despite resolution of the sepsis after anti-TB chemotherapy, she developed progressive ischaemia and impending gangrene of her fingers, toes, and feet, despite regular infusions of iloprost for 2 weeks. One day after starting sildenafil 50 mg




Figure 1 (A), (B), and (C) refer to cases 1, 2, and 3, respectively. In all three cases, ischaemia of viable tissues resolved after treatment with sildenafil (and tadalafil in case 2). (B) shows that increased finger extension became possible after treatment.



three times a day, all four extremities showed marked symptomatic improvement, becoming warm, less painful, and less discoloured. Her digital vasculitis resolved; weeks later necrotic zones became demarcated and autoamputated. Owing to resurgence of active lupus, her prednisolone dosage was increased; her haematological/serological abnormalities regressed. After 48 days of sildenafil treatment, tadalafil (Cialis) 10 mg once daily was substituted for 78 more days, without any adverse reaction. Currently, the patient remains well and can stand.

A 76 year old woman with prior hypertension and ischaemic heart disease (case 3) presented with erythematous facial and finger rashes, myalgia, finger and toe ischaemic lesions, and proximal muscle weakness. Dermatomyositis and a disseminated malignancy were diagnosed. After mutually agreeing to stop further investigation, prednisolone treatment was started. Her digital ischaemia

responded poorly to diltiazem SR and any improvement during iloprost infusions disappeared when they stopped. After 12 days and despite her past medical history, sildenafil treatment was cautiously started, with immediate reduction in peripheral pain and ischaemia and healing of toe ulcers. Although well tolerated, sildenafil was discontinued after 3 weeks, as the patient required treatment with nitrates.

DISCUSSION

All three patients had immediate, subjective and objective improvements after initiating sildenafil, which was entirely consistent with Lichtenstein's observations, except that our patients were very severely afflicted.⁴ In two patients, worsening RP and digital ischaemia appeared during active TB, consistent with the effects of sepsis.⁵ Rifampicin, a powerful inducer of relevant P450 enzymes (apt to enhance sildenafil elimination^{2,3}), appeared to affect their management. These

observations merit a randomised controlled trial to investigate the treatment of disabling RP with oral PDE-5 inhibitors.

.....
Authors' affiliations

C R Kumana, G T Y Cheung, C S Lau, Department of Medicine, The University of Hong Kong, 8/F Administration Block, Queen Mary Hospital, Hong Kong

Correspondence to: Professor C S Lau; cslau@hkucc.hku.hk

Accepted 29 December 2003

REFERENCES

- 1 **Michelakis ED**. The role of the NO axis and its therapeutic implications in pulmonary arterial hypertension. *Heart Fail Rev* 2003;**8**:5-21.
- 2 **Thummel KE**, Wilkinson GR. In vitro and in vivo drug interactions involving human CYP3A. *Annu Rev Pharmacol Toxicol* 1998;**38**:389-430.
- 3 **Warrington JS**, Shader RI, Von Moltke LL, Greenblatt DJ. In vitro biotransformation of sildenafil (Viagra): identification of human cytochromes and potential drug interactions. *Drug Metab Dispos* 2000;**28**:392-7.
- 4 **Lichtenstein JR**. Use of sildenafil citrate in Raynaud's phenomenon: comment on the article by Thompson et al. *Arthritis Rheum* 2003;**48**:282-83.
- 5 **Haj MA**, Robbie LA, Adey GD, Bennett B. Inhibitors of plasminogen activator in neutrophils and mononuclear cells from septic patients. *Thromb Haemost* 1995;**74**:1528-32.

Hereditary C1q deficiency and secondary Sjögren's syndrome

E P A H Hoppenreijns, P J van Dijken, P J Kabel, J M Th Draaisma

.....

Ann Rheum Dis 2004;**63**:1524-1525. doi: 10.1136/ard.2003.016592

A 13 year old Turkish boy with a known C1q deficiency and SLE-like disease developed recurrent parotitis. Investigations confirmed a secondary Sjögren's syndrome (SS). As far as we know, the association between C1q deficiency and SS has not been described before. The potential role of C1q in the pathogenesis of SS and systemic lupus erythematosus (SLE) might stimulate further research in understanding the pathogenesis of these and other autoimmune diseases. Screening patients with SS for complement deficiencies, including C1q deficiency, seems indicated.

Deficiency of the complement component C1q is a rare genetic disorder with susceptibility to recurrent infections with polysaccharide encapsulated micro-organisms and a high prevalence of autoimmune phenomena, most often SLE.

CASE REPORT

At the age of 4 years a boy of consanguineous Turkish descent had meningitis of unknown origin. At the age of 8 years he presented with meningococcal sepsis and meningitis. In the late convalescent phase of this infection he developed arthritis of his right elbow and pericarditis. At the age of 10 years he was admitted owing to lobular pneumonia. Immunological studies at that time demonstrated no functional or antigenic activity of C1q based on a homozygous Glu-86 stop mutation in the C1qA gene exon 2. Both parents and a sibling sister were found to be asymptomatic heterozygous carriers. His sibling brother, who also developed SLE-like symptoms, was homozygous for this mutation.

At the age of 13 years the patient developed recurrent arthritis of the ankle and elbow. Two years later he presented with six episodes of alternate right and left parotitis. There were no complaints of dry mouth or dry eyes. Ultrasonography of the parotid gland during an episode of parotitis disclosed diffuse swelling. Salivary scintigraphy showed delayed tracer uptake. Ocular examination showed some corneal erosion, a normal Schirmer test, and a tear break up time of 10 seconds. A salivary gland biopsy showed lymphocytic sialoadenitis with a focus score of 4 and a percentage of plasma cells containing IgA of 9%, consistent with SS. Microscopic haematuria and proteinuria were not present. Immunological studies showed positive antinuclear antibodies (ANA), RNP antibodies, anti-Sm antibodies, anti-SSA antibodies, and a positive rheumatoid factor (RF). Anti-SSB antibodies and anti-dsDNA antibodies were not

detected. A direct Coombs test and a serological test for syphilis were negative. Serum immunoglobulins were slightly raised. C3 and C4 were normal.

DISCUSSION

In our patient a homozygous point mutation in the C1qA gene was demonstrated that has earlier been described in five families from the Slovak republic and Turkey.¹ With arthritis, a positive ANA test, and anti-Sm antibodies our patient has an SLE-like disease.² Patients with SLE commonly have sicca symptoms, which may be related to the concomitant occurrence of SS.³ Secondary SS, in which the disease coexists with an autoimmune disease, is defined by the presence of either ocular or oral symptoms and two of four objective classification criteria.⁴ Our patient with recurrent swollen salivary glands, autoantibodies to ANA, RF, and SSA, a salivary scintigraphy showing delayed uptake, and biopsy findings consistent with SS, meets the full criteria for secondary SS.

These observations suggest that absence or abnormal function of C1q leads to susceptibility for SLE and SS. This may be due to ineffective immune complex clearance that causes tissue injury, exposes autoantigens, and stimulates an autoantibody response.⁵ In this respect it is significant that our patient had late onset reactive arthritis and pericarditis related to his meningococcal disease, which is also mediated by immune complexes.⁶ Other data suggest a defective clearance of apoptotic cells, which promotes accumulation of nucleosomes (suggested antigens in SLE), which in turn drives an autoimmune response.⁷

In conclusion we report a patient with C1q deficiency, SLE-like disease, and secondary SS. This is the first time that an association between C1q deficiency and SS has been reported. It suggests a role for C1q in the pathogenesis of autoimmune diseases, possibly due to ineffective immune complex clearing or defective apoptosis, and warrants further research.

Screening of patients with C1q deficiency for SS, but especially screening of patients with SS for complement deficiencies, including C1q deficiency, seems indicated.

.....
Authors' affiliations

E P A H Hoppenreijns, Department of Paediatrics, University Medical Centre Utrecht, The Netherlands

P J van Dijken, Department of Paediatrics, St Elisabeth Hospital Tilburg, The Netherlands

P J Kabel, Department of Medical Microbiology, St Elisabeth Hospital Tilburg, The Netherlands

J M Th Draaisma, Department of Paediatrics, University Medical Centre Nijmegen, The Netherlands

Correspondence to: Dr E P A H Hoppenreijns, Department of Paediatrics, University Medical Centre Utrecht, Wilhelmina Kinderziekenhuis, PO Box 85090, 3508 AB Utrecht, the Netherlands; e.p.a.h.hoppenreijns@wzk.azu.nl

Accepted 22 December 2003

REFERENCES

- 1 **Petry F**, Berkel AI, Loos M. Multiple identification of a particular type of hereditary C1q deficiency in the Turkish population: review of the

cases and additional genetic and functional analysis. *Hum Genet* 1997;**100**:51–6.

- 2 **Tan EM**, Cohen AS, Fries JF, Masi AT, McShane DJ, Rothfield NF, *et al*. The 1982 revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1982;**25**:1271–7.
- 3 **Gilboe IM**, Kvien TK, Uhlig T, Husby G. Sicca symptoms and secondary Sjögren's syndrome in systemic lupus erythematosus: comparison with rheumatoid arthritis and correlation with disease variables. *Ann Rheum Dis* 2001;**60**:1103–9.
- 4 **Viali C**, Bombardieri S, Jonsson R, Moutsopoulos HM, Alexander EL, Carsons SE, *et al*. European Study Group on Classification Criteria for Sjögren's Syndrome. *Ann Rheum Dis* 2002;**61**:554–8.
- 5 **Walport MJ**, Davies KA, Botto M. C1q and systemic lupus erythematosus. *Immunobiology* 1998;**199**:265–85.
- 6 **Finkelstein Y**, Adler Y, Nussinovitch M, Varsano I, Amir J. A new classification for pericarditis associated with meningococcal infection. *Eur J Pediatr* 1997;**156**:585–8.
- 7 **Bach JF**, Koutouzov S. Immunology: new clues to systemic lupus. *Lancet* 1997;**350**(suppl III):11.

Acute pneumonitis starting 2 hours after intramuscular gold administration in a patient with rheumatoid arthritis

A Hafejee, M J Burke

Ann Rheum Dis 2004;**63**:1525–1526. doi: 10.1136/ard.2003.017509

Gold salts have been used for the treatment of rheumatoid arthritis (RA) for over 50 years.¹ Their use is limited by the incidence of adverse reactions, including pneumonitis. Most case reports describe pneumonitis developing after a cumulative dose of several hundred milligrams.² Only four cases of lung disease have been reported with a cumulative dose of <100 mg.³

Here we report a case of gold pneumonitis, starting only 2 hours after the first maintenance dose of 20 mg was administered.

CASE REPORT

A 54 year old woman with RA had been followed up since 1984, her arthritis being well controlled with chloroquine. In 1997 her disease had flared up, and alternative disease modifying drugs were introduced—sulfasalazine, then methotrexate (both of were stopped because of adverse side effects), and finally, leflunomide. The last of these caused leucopenia and hypertension (controlled with bendrofluzide and ramipril) and hence was discontinued. Subsequently, her disease became more active again with prolonged joint stiffness, hand synovitis, and erythrocyte sedimentation rate (ESR) 102 mm/1st h.

Treatment was started with gold injections, with a test dose of 10 mg (no adverse effects noted), followed a week later by a dose of 20 mg. About 2 hours afterwards she complained of dizziness, nausea, then she developed a cough and breathlessness. She was admitted to hospital later that day. On examination she was afebrile, tachypnoeic, cyanosed (oxygen saturation 79% on air), pulse 110/minute, blood pressure 80/46, heart sounds normal, and had no signs of heart failure.

Investigations showed haemoglobin of 103 g/l with normal white cell and platelet counts. C reactive protein was 115 mg/l and troponin-I (a marker for myocardial infarction) was normal at 0.1 µg/l. Arterial blood gases on air showed a type I respiratory failure pattern. A chest x ray examination showed bilateral lower zone alveolar shadowing. An echocardiogram showed a normal ejection fraction.

After high dose oxygen treatment, intravenous hydrocortisone 100 mg, and intramuscular adrenaline 0.5 mg 1:1000, some improvement was seen clinically, oxygen saturation rising to 90% and blood pressure to 116/69. She was admitted to the high dependency unit and treatment started with pulse methylprednisolone 1 g daily, nasal continuous positive airway pressure ventilation, and required parenteral nutrition. On the day after her admission, she developed a generalised macular rash. Over the next 7 days her clinical condition improved (serial chest x ray examinations over this period showed clearing of the previous alveolar shadowing), and her treatment was changed to a reducing regimen of oral steroids. She was discharged, continuing frumil once daily and prednisolone 5 mg daily. Her arthritis was inactive and blood pressure 110/70. The ESR was 22 mm/1st h.

During her admission, it was considered that owing to the severity of her illness she was not fit enough to have other investigations—for example, computed tomography chest scan, bronchoalveolar lavage, and pulmonary function tests.

The gold was discontinued and her arthritis remains in remission.

DISCUSSION

Gold pneumonitis is a potentially fatal complication of gold salt treatment. This case illustrates two important points in management:

- Some of the early symptoms could have been confused with a nitritoid reaction. Indeed, hypertensive patients with RA who are taking angiotensin converting enzyme inhibitors (in this case ramipril) show a higher frequency of such reactions⁴ even if long term gold treatment is established. This combination should be avoided where possible.
- The clinical picture started to develop after only 2 hours of gold administration with just a 30 mg cumulative gold dose. Most case reports have suggested higher cumulative dose exposure^{2 5 6} and longer duration of treatment. To our knowledge such a florid onset has not been previously reported.